

Comparative Evaluation of Methanol Exposure and *Eucalyptus globulus* Extracts on Reproductive Parameters in Rats

HAMZA Dhikra¹, HAMDI Leila¹, ARKOUB Fatma Zohra², BOUKARINE Rahma², SAILI Linda³, KHELILI Kamel¹

¹Animal Ecophysiology Laboratory: Biology Department, Faculty of Science, Badji Mokhtar-Annaba University, Annaba 23000, Algeria.

²Laboratory of Environmental Sciences and Agro-ecology departement of biology, faculty of life and natural sciences, Chadli bendjedid El Tarf university, El Taref 36000, Algeria


³Environmental Research Center (CRE) ,Annaba , Algeria.

Email : dikra.hamza.dh@gmail.com

Abstract:

Plant-based biopesticides are considered as safer alternatives than synthetic pesticides. *Eucalyptus globulus* is widely recognized for its antioxidant potential, moreover the exposure to methanol raises the risk of possible reproductive toxicity. This study evaluated the effects of methanol and *Eucalyptus globulus* extracts on testicular function of rats. Forty rats were divided into four groups: Control, Methanol (600 mg/kg BW), Methanolic extract (ME, 100 mg/kg BW), and Aqueous extract (AE, 100 mg/kg BW). Treatments were administered orally for 28 days. At the end of the experiment, rats were sacrificed and testes and epididymides were collected. Sperm concentration, motility, and velocity were assessed using a CASA system. Serum testosterone levels and testicular oxidative stress markers (GSH, GPx, and MDA) were determined and histological evaluation of testicular tissue was performed using H&E staining. Methanol exposure was associated with the increase of oxidative stress, seminiferous tubule degeneration, and the reduce of sperm concentration, motility, and velocity. In addition , , rats treated with *Eucalyptus globulus* extracts showed more favorable antioxidant status, testicular histology, and sperm characteristics. In conclusion, methanol considerably affected testicular function and reproductive parameters, whereas *Eucalyptus globulus* extracts were associated with more favourable reproductive and oxidative stress profiles.

Keywords: Methanol, Eucalyptus extract, Oxidative stress, Reproductive health, Biopesticide.



Introduction:

Plant-based biopesticides are increasingly promoted as safer and more sustainable alternatives to synthetic pesticides, whose common use has been linked to environmental pollution and reproductive toxicity in humans and animals (Mnif et al., 2011; Mostafalou and Abdollahi, 2013). Among these natural sources, *Eucalyptus* species are particularly valued for their strong antioxidant and anti-inflammatory properties, which may associated with reduced oxidative stress–related damage in reproductive tissues (Djerrou et al., 2014; El-Sawi and Motawae, 2011).

To obtain these bioactive compounds, methanol is frequently used as an efficient organic solvent in extraction procedures. The methanolic extract, free from residual solvent, retains strong antioxidant and anti-inflammatory activities typical of the plant, thanks to its richness in phenolic compounds and flavonoids (Hariram and Soo, 2018). These properties may help counteract oxidative stress–mediated reproductive toxicity, offering potential protective effects on male fertility. Indeed, previous in vivo and in vitro studies have demonstrated the ability of *Eucalyptus* methanolic extracts to reduce oxidative damage in cells through scavenging of free radicals and enhancing antioxidant enzyme activity (Gonzalez-Burgos et al., 2018).

However, methanol itself is well known for its toxicity, especially its adverse effects on the male reproductive system, including impaired spermatogenesis, hormonal imbalance, and reduced fertility (Lee et al., 2010; Madsen and Larsen, 2012). These harmful effects are largely mediated by oxidative stress, characterized by excessive reactive oxygen species (ROS), lipid peroxidation, DNA damage, and germ cell degeneration (Agarwal et al., 2006; Tremellen, 2008).

.

Although the toxic effects of methanol were reported in several studies, its impact on male reproductive function remains insufficiently investigated. Furthermore, the limitation of the available information regarding the influence of *Eucalyptus globulus* extract on methanol-induced alterations in male reproductive function. Therefore, the present study aimed to evaluate the methanol effects on reproductive parameters and compare these findings with the *Eucalyptus globulus* effects extract administrated in male rats. The novelty of this study is based on the comparative examination of reproductive parameters, oxidative stress biomarkers, testosterone levels and histopathological changes to provide the evaluation of methanol-induced reproductive toxicity and the effects of *Eucalyptus globulus* extract.

Materials and methods:

1. Animals materials:

This study was conducted on 40 adult male Wistar rats with a mean body weight of 228.2 ± 20 g. The animals were obtained from the Pasteur Institute of Algeria and housed in groups of five per cage, with food and water provided *ad libitum*. They were put at the animal facility of Badji Mokhtar Annaba University, where they were allocated a 6-week acclimatization period under standard conditions. At the end of the experiment, the animals were anesthetized and then euthanized by decapitation in accordance with ethical guidelines.

2. Chemical material :

The methanol used in this experiment is purity $\geq 99\%$, which is a hydroxyl-group compound also known as methyl alcohol or carbinol, with the chemical formula CH_4O , purchased from (BIOCHEM Chemo-pharma, Montreal Quebec).

3. Plant material and preparation of aqueous extract:

3.1 Plant material:

Fresh *Eucalyptus globulus* leaves were obtained from the University of Badji Mokhtar, Annaba, in early 2023. The species was provided using standard morphological references, however a formal voucher specimen was not performed. After collection, the leaves were washed and dried in the shade, then ground into a coarse powder as part of the standard preparation process.

Aqueous extract :

The aqueous extract of *Eucalyptus globulus* was made by decocting method. The dried powdered leaf material was combined with distilled water at a 1:10 (w/v) ratio and brought to a boil at 100°C for 10 minutes. The mixture was then exposed to room temperature and filtered through Whatman No 1 filter paper and the filtrate was stored at temperature degree 4°C until further use.

3.2 Methanolic extract :

Dried and powdered *Eucalyptus globulus* leaves were cold-macerated in methanol at a solid-to-solvent ratio of 1:10 (w/v) for 24 hours in a sealed container at room temperature with regular stirring. The resulting mixture was then filtered through Whatman No 1 filter paper and concentrated under reduced pressure using a rotary evaporator at 40°C . The obtained extract was dried and stored at 4°C until further use.

4. The experimental design

After the adaptation period, the animals were divided into 4 groups (n=10) according to their body weight, ensuring that the mean body weight of each group was equal, as follows:

Group C (Control): received tap water;

Group M: received 600 mg/kg bw methanol dissolved in distilled water;

Group ME: received 100 mg/kg bw *Eucalyptus globulus* methanolic extract.

Group AE: received 100 mg/kg bw *Eucalyptus globulus* aqueous extract;

At the end of the-28 days of oral gavage treatment, all rats were sacrificed decapitation in accordance with ethical approval and institutional guidelines.

Sample collection:

5.1Hormone parameter assay (Testosterone):

Plasma testosterone concentration was determined using a competitive ELISA kit (DRG EIA-1559, DRG Instruments GmbH, Germany), following the manufacturer's instructions. The competitive ELISA is based on competition between sample antigen and enzyme-labeled antigen for antibody binding sites in microplate wells. After incubation (1–2 h) and washing, TMB substrate was added to produce a blue color, which was stopped with 1N HCl, turning yellow. Absorbance was read at 450 nm. All samples were analyzed in duplicate, with an intra-assay CV <10%.

5.2 Study of the biology of spermatozoa:

Sperm analysis was carried out using the SCA® CASA system (Sperm Class Analyzer® Computer-Assisted Sperm Analysis). In order to assess sperm concentration and motility, samples were collected from a small incision in the cauda epididymis. After a 1:8 dilution of semen in 0.9% NaCl physiological solution, a drop was transferred using a micropipette to a GoldCyto 20-µm counting chamber. The preparation was then examined under a microscope (Nikon Eclipse E200) maintained at 37 °C, using ×4 negative-phase contrast combined with a phase-contrast condenser and integrated into the microcomputer.

5.3 Oxidative stress assays:

After dissection, the testes were removed, carefully cleaned of fatty tissue, and weighed using a precision balance (Kern PRS 320-3). The tissues were then homogenized in cold phosphate buffer (0.1 M, pH 7.4) using a homogenizer at 4°C. They were then used to determine:

- **Malondialdehyde (MDA) levels** following the method of Ohkawa *et al.*, (1979).
- **Reduced glutathione (GSH) levels** according to Weekbeker and Cory (1988).
- **Glutathione peroxidase (GPx) activity** based on the method of Flohe and Günzler (1984).
- **Total protein concentration** using Bradford's method (1976).

The results were expressed as following:

- nmol/mg tissue for MDA.
- µM GSH/mg protein for GPx activity and GSH.

5. Histopathological study

Formaldehyde-fixed samples were dehydrated through a graded series of alcohols (70% to 100%) using a circulating tissue processor (Thermo Scientific, Microm STP 120, UK) and subsequently embedded in paraffin wax. Paraffin sections of 5 µm thickness were mounted on clean glass slides, stained with hematoxylin and eosin (H&E) (BioLab, Mumbai, India), and examined under a light microscopic at ×40 magnification (Leica DMLB, Wetzlar, Germany) (Hould, 1984).

Statistical analysis:

Statistical significance was performed using GraphPad Prism 9 (GraphPad Software). Statistical significance was determined using a one-way analysis of variance (ANOVA) followed by Tukey's post-hoc test. Data are expressed as mean ± Standard Deviation (mean ± SD). Differences was considered significant at $p < 0.05$.

Results:

1. Reproductive profile:

1.1 Testes and epididymis weight:

The absolute weights of the reproductive organs (testes and epididymis) are presented in Table 01. Exposure to methanol (M) decreased the testicular weight compared with the control group ($p < 0.05$). Similarly, epididymal weight was lower in the methanol-treated group (M) compared with the control group ($p < 0.05$). On the other hand, treatment with the methanolic extracts of *Eucalyptus globulus* (ME) showed higher testicular and epididymal weights compared with the methanol group (M).

Table 1: Absolute weights of testes and epididymis in the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus* extract; AE: Aqueous *Eucalyptus* extract.

Groups	Testes weight	Epididymis weight
C	1.850 $\pm$ 0,3119 ^a	0,6315 $\pm$ 0,08520 ^a
M	1,581 $\pm$ 0,1093 ^b	0,3979 $\pm$ 0,07878 ^b
ME	1,878 $\pm$ 0,1733 ^a	0.6707 $\pm$ 0,04255 ^a
AE	1,738 $\pm$ 0,1075 ^{a.b}	0,6060 $\pm$ 0,07001 ^a

1.2 Sperm Concentration:

As shown in figure 1, sperm concentration was reduced in the methanol-treated group in comparison to the control group ($p < 0.05$). Treatment with the eucalyptus methanolic extract (ME) and the aqueous extract (AE) showed a higher sperm concentration compared to the methanol (M) group ($p < 0.05$).




Figure 1: Sperm concentration in the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus* extract; AE: Aqueous *Eucalyptus* extract.

1.3 Sperm motility:

Figure 2 shows that sperm motility was reduced in the methanol-treated group compared with the control group ($p < 0.05$).




Figure 2: Total sperm motility in rats from the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus* extract; AE: Aqueous *Eucalyptus* extract.

1.4 Sperm velocity:

Across the fast sperm category, as presented in figure 3, no differences were observed among all experimental groups ($p > 0.05$). In the medium-velocity category, values were lower in the methanol group (M) compared with the control ($p < 0.05$). Similarly, slow-moving sperm showed a decrease in the methanol group ($p < 0.05$). In contrast, the proportion of immotile sperm was higher in the methanol-treated group compared with the control ($p < 0.05$). Importantly, both (ME) and (AE) groups showed no differences from the control across all sperm velocity categories ($p > 0.05$).




Figure 3: Sperm velocity in rats across the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus globulus* extract; AE: Aqueous *Eucalyptus globulus* extract.

2. Testosterone levels:

The testosterone levels (Figure 4) were lower in the methanol (M) group compared with the control group ($p < 0.05$). Treatment with the methanolic (ME) and aqueous (AE) extracts of *Eucalyptus globulus* showed higher testosterone levels compared with the methanol (M) group ($p < 0.05$).




Figure 4: Testosterone levels in rats across the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus globulus* extract; AE: Aqueous *Eucalyptus globulus* extract.

3. Oxidative stress markers:

3.1 Malondialdehyde (MDA) levels:

The MDA levels (Figure 5) were higher in the methanol (M) group compared with the control group ($p < 0.05$). In contrast, both the ME and AE groups showed no significant differences compared with the control group ($p > 0.05$).




Figure 5: Testicular malondialdehyde (MDA) levels in rats across the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus globulus* extract; AE: Aqueous *Eucalyptus globulus* extract.

3.2 Reduced Glutathione (GSH) content:

GSH levels (Figure 8) were reduced in the methanol (M) group compared with the control group. No significant differences were observed in the ME and AE groups relative to controls ($p > 0.05$).




Figure 6: Testicular reduced glutathione (GSH) content in rats across the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus globulus* extract; AE: Aqueous *Eucalyptus globulus* extract.

3.3 Peroxidase Glutathione (GPx) activity:

Similarly, GPx levels (Figure 7) were reduced in the methanol (M) group compared with the control group. Conversely, higher GPx levels were observed in the groups treated with the aqueous eucalyptus extract (E) and the methanolic eucalyptus extract (ME) compared with the control (C) group ($p < 0.05$).




Figure 7: Testicular glutathione peroxidase (GPx) activity in rats across the different experimental groups (mean $\pm$ SD). Values that do not share the same superscript letters are significantly different at $p < 0.05$. C: Control; M: Methanol; ME: Methanolic *Eucalyptus globulus* extract; AE: Aqueous *Eucalyptus globulus* extract.

4. Histological study:

4.1 Testes:

Microscopic examination of testicular sections in the **control group** revealed normal histo-architecture, with well-organized seminiferous tubules, an intact germinal epithelium and numerous spermatozoa in the lumen. In the **methanol** treated group, marked histological alterations were observed, including **empty lumina**, **vacuolization of Sertoli cells**, and **atrophy of seminiferous tubules**, reflecting severe disruption of spermatogenesis and

testicular degeneration. In contrast, rats treated with the **methanolic (ME) and aqueous (AE)** **extracts of *Eucalyptus globulus*** showed nearly normal testicular architecture, with preserved germinal layers and evidence of active spermatogenesis, suggesting a protective effect of both extracts.




Figure (08): Microscopic observation of histological sections of the testes. (A) Control group;

(B) M group; (C) ME group; (D) AE group. H&E staining, magnification $\times 10$ (left) and $\times 40$ (right). ST: Seminiferous tubule, TL: Tubular lumen, IT: Interstitial tissue, LC: Leydig cell, SC: Sertoli cell, SPZ: Spermatozoa, EL: Empty Lumen, E: interstitial edema, V: vacuolization in Sertoli cells. Atrophy of the seminiferous tubules (Black arrow).

4.2 Epididymis:

Microscopic examination of epididymal sections in the **control group** revealed a normal histoarchitecture, with well-defined epididymal ducts and lumina densely filled with mature spermatozoa. Conversely, the **methanol-treated group** displayed marked histopathological alterations characterized by **empty lumina** and a **notable reduction in sperm count**, reflecting impaired sperm storage and transport, consistent with methanol-induced reproductive toxicity. In rats treated with **methanolic (ME) and aqueous (AE) extracts of *Eucalyptus globulus***, the epididymal epithelium appeared preserved, and the ducts remained rich in spermatozoa, indicating maintained epididymal function and sperm maturation.




Figure (09): Microscopic observation of histological sections of the epididymis. (A) Control group; (B) Methanol (M) group; (C) Aqueous eucalyptus extract (E) group; (D) Methanolic eucalyptus extract (ME) group. H&E staining, magnification $\times 10$ (left) and $\times 40$ (right).

EC: Epididymal canal, LEC: Lumen of the epididymal canal, EL: Empty lumen, BM: Basement membrane, SPZ: Spermatozoa. (↓): Decrease in sperm count.

Discussion:

Reproductive disorders refer to a range of conditions that impair the ability of the reproductive system to function properly. Numerous studies have shown that exposure to environmental contaminants, especially pesticides, negatively affects the reproductive health of both males and females (Kumar, 2004; Shojaei Saadi and Abdollahi, 2012). International studies have confirmed the reproductive toxicity of various solvents and have identified the testes as primary target organs for toxic substances released into the environment (Fort et al., 2001).

Our findings showed that exposure to 600 mg/kg body weight of methanol over a 28-day period was associated with reduced sperm concentration, motility, and velocity. These results are in agreement with those reported by Tielemans et al., who observed impaired sperm quality in workers exposed to solvents. Furthermore, several studies have provided experimental evidence of additional impairments in the spermatogenesis process, including disturbances in sperm cell division, reduced motility, and alterations in sperm density and viscosity. Evidence suggests that male fertility may be adversely affected by various toxicants, either through direct damage to testicular tissue or by disrupting the endocrine regulation of male reproductive function (Jeng, 2014). Notably, even short-term exposure may be sufficient to disturb the hormonal balance required for normal spermatogenesis, as reflected by changes in testosterone and other hormone levels. These findings suggest that methanol exposure may affect both hormonal regulation and sperm production in male rats (Cooper et al., 1992).

Methanol intoxication has been associated with mitochondrial dysfunction and increased microsomal enzyme proliferation, both of which contribute to the overproduction of reactive oxygen species (ROS) (Tephly, 1991). The observed reduction in glutathione (GSH) levels may also explain the decreased concentration of vitamin C, as vitamin C is primarily absorbed by cells in its oxidized form and subsequently regenerated by GSH (Briviba and Sies, 1994). Oxidative stress is defined as a significant imbalance between the production of oxidizing agents and the organism's antioxidant defense capacity. This imbalance arises either from excessive generation of reactive oxygen species or from impairments in the antioxidant defense system (Dahmoune et al., 2015; Boyd et al., 2003).

The present study revealed that both the aqueous and methanolic extracts of *Eucalyptus globulus* were associated with increase in GSH and GPx levels, along with decrease in MDA levels in the testes. Glutathione (GSH) plays a vital role in the cellular defense system against toxic compounds of both endogenous and exogenous origin (Meister and Anderson, 1983). Beyond its well-known antioxidant properties, GSH is also involved in protein and DNA synthesis, maintenance of cell membrane integrity, and regulation of enzymatic activities (Meister, 1989). Studies have demonstrated that the enzyme glutathione peroxidase, which functions as an antioxidant, plays a critical role in protecting sperm within testicular and epididymal tissues. A deficiency of this enzyme has been linked to infertility. Located in the plasma membrane of sperm, the sperm nucleus, and epididymal fluid, glutathione peroxidase protects sperm from free radical damage and contributes to the final maturation and development of spermatozoa (Chabory et al., 2009; Drevet, 2006). Moreover, malondialdehyde (MDA), a byproduct of lipid peroxidation, is widely used as a reliable biomarker to assess the level of oxidative stress in biological systems. Elevated levels of MDA indicate increased cellular membrane damage due to reactive oxygen species (ROS), which can contribute to various pathological conditions, including reproductive toxicity (Xiong et al., 2016).

The three tested extracts exhibited significant antioxidant properties, with their potency varying according to the in vitro assay employed. Notably, *Eucalyptus globulus* extracts effectively mitigated H₂O₂-induced oxidative stress by enhancing cell viability, increasing glutathione (GSH) levels and antioxidant enzyme activities, while concurrently reducing reactive oxygen species (ROS) production and lipid peroxidation in SH-SY5Y cells (Tremellen, 2008). Interestingly, *Eucalyptus globulus* mitigated diclofenac sodium-induced alterations in reproductive parameters by enhancing the cellular antioxidant system, leading to a significant increase in glutathione (GSH) concentration within testicular tissue (Mousa et al., 2019). *Eucalyptus globulus* leaf extracts may serve as valuable raw materials for food and pharmaceutical supplements due to their high content of antioxidant compounds, which confer protective health benefits against oxidative stress (Tremellen, 2008).

Eucalyptus leaves are considered a source of antioxidants, particularly flavonoids, which may contribute to protection against oxidative stress and damage caused by free radicals (Nakhaee et al., 2009).

This study shows that administration of the methanolic extract of *Eucalyptus globulus* was associated with alterations in serum total testosterone levels, as well as improvements in sperm

parameters, including sperm count and individual motility. These enhancements are likely attributed to the antioxidant properties of *Eucalyptus*. We hypothesize that *Eucalyptus* supplementation may stimulate the production of reduced glutathione in testicular tissues, thereby protecting spermatozoa from oxidative damage and preserving their motility (Mousa et al., 2019). An increase in testicular weight may occur due to elevated testosterone secretion, stimulation of germ cell proliferation and division, as well as the effects of terpenes and flavonoids present in the extract (Yakubu, 2012) .

Today, there is growing interest in the use of biopesticides as a more sustainable and reliable alternative to synthetic chemical pesticides. Biopesticides are derived from biological sources, including living organisms or substances extracted from natural origins. More broadly, the term encompasses any plant protection product that is not based on synthetic chemical compounds (Yovo, 2010).

Recent studies have demonstrated that exposure to endocrine-disrupting chemicals, including pesticides, mercury, phenols, and phthalates, can interfere with hormone metabolism by altering the synthesis and/or degradation of testosterone, follicle-stimulating hormone (FSH), luteinizing hormone (LH), and other key reproductive hormones. Steroid metabolism primarily occurs in the liver, which is also a major target for many exogenous toxicants. Although numerous scientific publications have explored this topic, a clear causal relationship between such hormonal disturbances and reproductive disorders such as infertility, azoospermia, or gonadal dysfunction has yet to be definitively established (Queiroz and Waissmann, 2006; Alejandro et al., 2002; Dalgaard et al., 2001).

Microscopic examination showed differences in histological features and spermatogenesis among the groups. In the control group and the *Eucalyptus globulus*-treated groups, the seminiferous tubules generally maintained a normal structure, with ongoing spermatogenesis and the presence of spermatozoa within the lumen. In contrast, methanol-treated rats showed noticeable changes in the seminiferous tubules, . With an altered appearance and a reduced presence of spermatozoa in the lumen. An increase in the intertubular space was also observed, along with vacuolization of Sertoli cells. In the epididymis a lower concentration of spermatozoa was observed in the methanol-treated group, together with structural alterations in the epididymal ducts.

The histoarchitecture of the testes in both the control and *Eucalyptus globulus*-treated groups exhibited compactly arranged seminiferous tubules containing germinal layers at all stages of spermatogenesis, Sertoli cells, a normal lumen filled with spermatozoa, intact interstitial tissue, and functional Leydig cells (Mousa et al., 2019)]. Sertoli cells play a critical role in supporting spermatogenesis (Sapori et al., 1986), while Leydig cells are the primary source of androgen production. Both cell types are susceptible to damage from toxicants and chemical drugs (Papadakis et al., 1999).

This study has several limitations that should be acknowledged. First, only a single methanol dose and a 28-day exposure period were evaluated; therefore, the dose-dependent and long-term effects of methanol remain to be investigated. In addition, although oxidative stress biomarkers, reproductive hormone levels, sperm parameters, and histopathological changes were assessed, the underlying molecular mechanisms responsible for the protective effects of *Eucalyptus globulus* were not explored. Future studies incorporating multiple dose regimens, longer exposure periods, and molecular analyses are warranted to further elucidate the mechanisms underlying the protective effects of *Eucalyptus globulus* against methanol-induced reproductive toxicity.

Conclusion:

This study demonstrates that *Eucalyptus globulus* extracts, particularly the methanolic and aqueous forms, are associated with male reproductive parameters improvement. Both extracts reduced oxidative stress markers, lowered lipid peroxidation, and improved antioxidant enzyme activities, reflecting the strong antioxidant and anti-inflammatory potential of *Eucalyptus* compounds. The slight increase in testosterone levels observed in treated groups suggests an effect on testicular function. However, the methanol exposure was associated with reproductive toxicity, characterized by level testosterone decrease, this exposure elevated oxidative stress and impaired testicular integrity.

Overall, these findings highlight differences between methanol exposure and *Eucalyptus globulus* extract treatment in reproductive parameters, and suggest that further investigation of these extracts may be a topic of interest. Further studies are required to clarify the above mechanisms and determine optimal dosing for practical use.

409 **References:**

- 410 Agarwal A, Gupta S, Sharma RK. 2006. Role of oxidative stress in female reproduction. *Reproductive*
411 *Biology and Endocrinology*. 4(1):49.
- 412 Alejandro A, Giami A, Multigner L. 2002. Environmental agents and erectile dysfunction: a study in a
413 consulting population. *Journal of Andrology*. 23:546–550.
- 414 Boyd B, Ford C, Koepke M, Gary K, Horn E. 2003. Étude pilote ouverte de l'effet antioxydant
415 d'Ambrotose sur des personnes en bonne santé. *Glycoscience Nutrition*. 4(6):70–77.
- 416 Briviba K, Sies H. 1994. Nonenzymatic antioxidant defense systems. In: Frei B, editor. *Natural*
417 *Antioxidants in Human Health and Disease*. San Diego: Academic Press. p. 107–128.
- 418 Chabory E, Damon C, Lenoir A, et al. 2009. Epididymis seleno-independent glutathione peroxidase 5
419 maintains sperm DNA integrity in mice. *Journal of Clinical Investigation*. 119(7):2074–2085.
- 420 Cooper RL, Mole ML, Rehnberg GL, Goldman JM, McElroy WK, Hein J, Stoker TE. 1992. Effect of
421 inhaled methanol on pituitary and testicular hormones in chamber-acclimated and non-acclimated rats.
422 *Toxicology*. 71(1–2):69–81.
- 423 Dalgaard M, Hossaini A, Hougaard KS, et al. 2001. Developmental toxicity of toluene in male rats:
424 effects on semen quality, testis morphology, and apoptotic neurodegeneration. *Archives of Toxicology*.
425 75:103–109.
- 426 Dahmoune F, Nayak B, Moussi K, Remini H, Madani K. 2015. Optimization of microwave-assisted
427 extraction of polyphenols from *Myrtus communis* L. leaves. *Food Chemistry*. 166:585–595.
- 428 Djerrou Z, Maameri Z, Hamdi-Pacha Y, et al. 2014. Antioxidant effect of *Eucalyptus globulus* against
429 lead-induced oxidative stress in reproductive organs in male rats. *Journal of Environmental Health*
430 *Science and Engineering*. 12:11.
- 431 Drevet JR. 2006. The antioxidant glutathione peroxidase family and spermatozoa: a complex story.
432 *Molecular and Cellular Endocrinology*. 250(1–2):70–79.
- 433 El-Sawi SA, Motawae HM. 2011. Antioxidant and anti-inflammatory activities of essential oils from
434 eucalyptus species. *Journal of Applied Pharmaceutical Science*. 1(6):33–37.
- 435 Fort DJ, Stover EL, Bantle JA, Dumont JN, Finch RA. 2001. Evaluation of a reproductive toxicity assay
436 using *Xenopus laevis*: boric acid, cadmium and ethylene glycol monomethyl ether. *Journal of Applied*
437 *Toxicology*. 21(1):41–52.
- 438 Gonzalez-Burgos E, Liaudanskas M, Viskelis J, Zvikas V, Janulis V, Gomez-Serranillos MP. 2018.
439 Antioxidant activity, neuroprotective properties and bioactive constituents analysis of varying polarity
440 extracts from *Eucalyptus globulus* leaves. *Journal of Food and Drug Analysis*. 26(4):1293–1302.
- 441 Hariram NS, Soo KY. 2018. Hydropenia induces expression of drought responsive genes in *Linum*
442 *usitatissimum* L. *Indian Journal of Experimental Biology*. 56(10):743–749.
- 443 Hould R. 1984. Techniques d'histopathologie et de cytopathologie. In: Decarie M, editor. Montreal:
444 Maloine. p. 225–227.
- 445 Jeng HA. 2014. Exposure to endocrine disrupting chemicals and male reproductive health. *Frontiers in*
446 *Public Health*. 2:55.

447 Kumar S. 2004. Occupational exposure associated with reproductive dysfunction. *Journal of*
448 *Occupational Health*. 46(1):1–19.

449 Lee HS, Kim SA, Park J. 2010. Reproductive toxicity of methanol and its metabolite, formic acid.
450 *Journal of Reproductive Toxicology*. 29(4):375–380.

451 Madsen AM, Larsen ST. 2012. Cytotoxic and inflammatory effects of methanol vapors on human airway
452 epithelial cells. *Toxicology Letters*. 213(2):278–284.

453 Meister A. 1989. Metabolism and function of glutathione. In: Dolphin D, Poulsen R, Avramovic O,
454 editors. *Glutathione: Chemical, Biological and Medical Aspects*. New York: Wiley. p. 367–374.

455 Meister A, Anderson S. 1983. Glutathione. *Annual Review of Biochemistry*. 52:711–760.

456 Mnif W, Hassine AIH, Bouaziz A, et al. 2011. Effect of endocrine disruptor pesticides: a review.
457 *International Journal of Environmental Research and Public Health*. 8(6):2265–2303.

458 Mostafalou S, Abdollahi M. 2013. Pesticides and human chronic diseases: evidences, mechanisms, and
459 perspectives. *Toxicology and Applied Pharmacology*. 268(2):157–177.

460 Mousa AA, Elweza AE, Elbaz HT, Tahoun EAA, Shoghy KM, Elsayed I, Hassan EB. 2019. *Eucalyptus*
461 *globulus* protects against diclofenac sodium-induced hepatorenal and testicular toxicity in male rats.
462 *Journal of Traditional and Complementary Medicine*.

463 Nakhaee A, Bokaeian M, Saravani M, Farhangi A, Akbarzadeh A. 2009. Attenuation of oxidative stress
464 in streptozotocin-induced diabetic rats by *Eucalyptus globulus*. *Indian Journal of Clinical Biochemistry*.
465 24(4):419–425.

466 Papadakis V, Vlachopapadopoulou W, Van Syckle K, et al. 1999. Gonadal function in young patients
467 successfully treated for Hodgkin's disease. *Medical and Pediatric Oncology*. 32(5):366–370.

468 Queiroz EKR, Waissmann W. 2006. Occupational exposure and effects on the male reproductive
469 system. *Cadernos de Saúde Pública*. 22:485–493.

470 Saporì PMH, Chatelain P, Saez JM. 1986. In vitro interaction between Sertoli cells and steroidogenic
471 cells. *Biochemical and Biophysical Research Communications*. 134(3):957–962.

472 Shojaei Saadi H, Abdollahi M. 2012. Is there a link between human infertilities and exposure to
473 pesticides? *International Journal of Pharmacology*. 8(8):708–710.

474 Tremellen K. 2008. Oxidative stress and male infertility—a clinical perspective. *Human Reproduction*
475 *Update*. 14(3):243–258.

476 Xiong F, Li Q, Zhou B, Huang J, Liang G, Zhang L, et al. 2016. Oxidative stress and genotoxicity of
477 BTEX exposure. *International Journal of Environmental Research and Public Health*. 13(12):1212.

478 Yakubu MT. 2012. Effect of *Chromolaena odorata* extract on reproductive indices. *Journal of*
479 *Andrology*. 33(6):1199–1207.

480 Yovo K. 2010. Consentement à payer les biopesticides. *Tropicultura*. 28(2):101–106.

481

482

483