

1 **Nanopriming with chitosan-coated SiO₂NPs alleviates cadmium toxicity in wheat (*Triticum***
2 ***aestivum* L.) seedlings: an ecofriendly and sustainable approach**

3 **Ayesha Saeed¹, Razia Noreen¹, Nosheen Aslam¹, Arruje Hameed^{1,*},**

4 ¹*Department of Biochemistry, Government College University Faisalabad, Pakistan*


5 **Full Postal Address:**

6 Department of Biochemistry,
7 Government College University Faisalabad,
8 Faisalabad, Punjab, Pakistan

9 ***Corresponding Author:**

10 Arruje Hameed

11 Email: arrujeh@yahoo.com, arrujehameed@gcuf.edu.pk



Abstract

Heavy metals have become a major threat to the yield and grain quality of economically important cereal crops. Cadmium (Cd) is highly toxic and disrupts key physiological functions, biochemical processes, and cellular structures, retarding germination and seedling development. In recent years, SiO₂NPs have shown promising applications in sustainable agriculture. In this study, chitosan (CS)-coated SiO₂NPs were prepared and employed as the nanopriming agents to mitigate toxic effects of Cd in wheat seedlings. Nanopriming with CS-coated SiO₂NPs readjusted the biochemical parameters, which counteracted Cd stress (5 µM) in wheat seedlings compared to the control. They significantly upregulated the hydrolytic enzymes (protease and α-amylase), and enzymatic antioxidants (CAT, POD and SOD). They also induced a significant increase in sugars (reducing sugars, non-reducing sugars, and total sugars) and total soluble proteins, alleviating the Cd toxicity in wheat seedlings. The activated antioxidant potential controlled ROS production, avoiding lipid peroxidation and reducing MDA content under Cd stress. The nanopriming-modulated biochemical attributes as a countermeasure of Cd toxicity, ensuring sustainable seedling development. The highest concentration of Cd (BT₃; 1.5 ppm) caused the greatest overall biochemical response, resulting in a 51.7%, 49.5%, 49.8%, and 33.5% increase in TSP, CAT, POD, and SOD, respectively, compared to the non-primed control, but decreasing MDA by 20.1%. The responses were generally more pronounced in BT₃ than the uncoated SiO₂NP treatment (AT₃), indicating the greater efficacy of chitosan coating on nanopriming response under Cd stress.

Keywords: Seed pre-conditioning, Climate-smart, Sustainable agriculture, Heavy metal toxicity, Cadmium stress

1. Introduction

Over the last few decades, heavy metals (HMs) have emerged as main soil pollutants that reduce yield of field crops, posing food security challenge and threatening the recent trends in sustainable agriculture. They are causing huge revenue loss to agro-based economies, especially in developing countries (Hou et al., 2020; Rashid et al., 2023). Cadmium (Cd) metal is frequently used in various

industrial processes, such as the production of paints, pesticides, plastics, textile dyes, and rechargeable batteries. Therefore, Cd is causing air, water and soil pollution, resulting in numerous negative impacts on the environment. It shows toxicity to germinating seeds, seedling development and plant growth, leading to low crop yield (Huybrechts et al., 2019). It causes ROS production, damaging the antioxidant machinery and sub-cellular signaling mechanisms (Hassan et al., 2020). Its toxic nature damages metabolism, the respiratory process and energy production (Zhang et al., 2020). It negatively influences the biosynthesis of proteins, sugars and secondary metabolites in germinating seeds and plants. It affects photosynthesis by disturbing the ionic balance and turgor pressure across the cells (Aslam et al., 2023; Liu et al., 2020). Such changes in biochemical attributes due to Cd toxicity retards germination and plant growth, which undermines qualitative and quantitative crop performance (El Rasafi et al., 2022). Wheat (*Triticum aestivum* L.) is a major staple food and economically valuable cereal crop around the globe, especially in Asia. Excessive Cd levels reduce its productivity, causing significant economic losses. Generally, plants upregulate antioxidants and growth hormones to counter Cd toxicity (Gill et al., 2021; Zhu et al., 2021). Recently, nanopriming has become an easy seed pre-conditioning technique that supports them to tolerate abiotic stresses, including the toxic HMs (Bhatia & Gupta, 2022; Kumar et al., 2021).

The primed seeds resist HM's toxicity through several countermeasures, including activation of antioxidants and metabolic activities, readjustment of osmolytes, and cell signaling (Wiszniewska, 2021; Wojtyla et al., 2016). In recent years, nanomaterials have received considerable attention in sustainable agriculture owing to their wide range of applications for managing stress and disease in plants. They have been widely explored as climate-smart priming agents against biotic and abiotic stresses (Sarkar et al., 2022; Singh et al., 2022). Over the last few years, SiO₂NPs have shown promising applications in plant growth regulation owing to their stable and non-toxic nature (Wang et al., 2022; Weisany et al., 2024). Chitosan (CS), a natural biopolymer, has become a promising material with broad applications in plant growth regulation. It positively influences physiological and biochemical attributes in growing plants. It can upregulate antioxidants and increase biosynthesis of

phytoalexin and other potent biomolecules (Mukarram et al., 2023). It increases osmolyte levels and stomatal conductance, improving photosynthetic activity. It can act as an antiviral, antibacterial and antifungal agent, thus showing potential applications for disease management in plants (Nandhini et al., 2025; Suwanchaikasem et al., 2024). The nanoformulation of CS, such as CSNPs, has become an attractive nano-carrier for controlled release of fertilizers, pesticides and other growth-promoting agents (Ma et al., 2023; Saberi Risch et al., 2024).

The application of SiO₂NPs and CS using foliar spray, soil application and conventional seed treatment has been previously well studied, whereas seed nanoprimering represents a different and more targeted mode of delivery (Khan et al., 2023; Nile et al., 2022). It provides active agents directly in the seed matrix in the pre-germination stage, allowing the metabolic reprogramming to occur at an early stage before exposure to Cd (Eevera et al., 2023; Khan et al., 2023). The combined nanoformulation of CS and SiO₂NPs was hypothesized to cumulate the beneficial effects of both the components, which may lead to synergistic antioxidant priming and better imbibition in the seed.

Considering the benefits of nanoprimering and promising applications of CS and SiO₂NPs, we envisioned preparing CS-coated SiO₂NPs as an ecofriendly nanoprimering agent to evaluate their efficacy against Cd toxicity in wheat seedlings.

2. Materials and Methods

2.1 Preparation of CS-coated SiO₂NPs as nanoprimering agent:

The 3g of Na₂SiO₃ was dissolved in 15 mL of distilled water and stirred at RT for 6 hr. Then, the solution was neutralized with 1 M HCl, subsequently filtered and dried at 65 °C to obtain the white powder of SiO₂NPs (sample A).

The 50 mg of SiO₂NPs were slowly dissolved in 10 mL of 1% CS solution ((w/v) in acetic acid) followed by sonication and stirred for 24 hr. at RT. Afterwards, the mixture was subjected to centrifugation, rinsed with deionized water, and dried at 65 °C to obtain CS-coated SiO₂NPs (sample B) (**Figure 1**) (Saleh et al., 2022).




Figure 1. Preparation of CS-coated SiO₂NPs as nanopriming agent

The SEM analysis revealed a layered mesoporous structure of SiO₂NPs and a spherical nature of clustered CS-coated SiO₂NPs (**Figure 2a, b**). SEM images were obtained from an FEI Nova NanoSEM 450 field emission scanning electron microscope that is fitted with a field-emission gun. Imaging was done at the accelerating voltage of 10 kV, spot size 2.5 and working distance of ~ 5.06 mm under high-vacuum condition, through the Through the Lense Detector (TLD) in secondary electron (SE/CN) signal mode.

The functional groups' interactions were detected using FTIR analysis (**Figure 2c**). FTIR spectra were collected on the Agilent Technologies MicroLab spectrometer with a spectral resolution of 4 cm⁻¹. A total of 30 sample scans and 10 background scans were recorded in the spectral range 4000-650 cm⁻¹, using the Happ-Genzel apodization function during spectral acquisition.

The XRD validated the crystalline nature of SiO₂NPs and the semi-crystalline nature of CS-coated SiO₂NPs (**Figure 2d**). The XRD patterns were collected with a Bruker D8 Advance diffractometer with a Theta-Theta goniometer and LynxEye position sensitive detector (PSD). The

radiation was Cu-K α with operating voltage and current of 30 kV and 25 mA, respectively. Continuous PSD fast-scan mode was used to collect data in the range 5° - 80° 2 θ and with a step size of about 0.02° and a step time of 0.025 s.




Figure 2. Characterization of CS-coated SiO₂NPs

2.2 Nanoprimering and germination of wheat seeds:

Spring wheat (*Triticum aestivum* L. cv. Akbar-2019) was obtained from The Ayub Agriculture Research Institute, Faisalabad, Pakistan for this study. The wheat seeds were primed using SiO₂NPs (sample A) and CS-coated SiO₂NPs (sample B). The 0.5 (T₁), 1 (T₂) and 1.5 (T₃) ppm solutions of A and B were prepared for nanoprimering of 25 g of seeds for 8 hrs at 23±2°C. Afterwards, they were dried to their original weight.

In parallel, some seeds were hydro-primed for comparison with nanoprimered and non-primed controls. The control, nanoprimered and hydroprimed seeds were allowed to germinate in 12 cm diameter petri plates at 23±2°C (24 seeds in three independent biological replicates (n=3)) (Anonymous, 1990) under Cd stress (5 μ M CdCl₂) and normal conditions (Ci et al., 2010). The germinating seeds were counted twice a day for 2 weeks. The germination data was recorded and used to determine the percentage of germination, germination index, energy of germination and mean

germination time. The entire germination data will be reported in the separate whole-plant study for whole-plant growth responses. The seedling samples were stored at -80 °C before biochemical analyses.

2.3 Biochemical studies of wheat seedling:

For antioxidant enzyme activity, seedling samples (0.1-0.2 g fresh weight) were homogenized in ice-cold 50 mM potassium phosphate buffer (pH 7.0 for CAT and POD; pH 7.8 for SOD; 1:10 w/v). For protein and sugar content, seedling samples (0.1-0.2 g fresh weight) were homogenized in 0.1% (w/v) TCA (1:10 w/v). Homogenates were centrifuged, at 10,000 x g for 20 mins at 4°C, and the supernatant was used for biochemical assays as described.

2.3.1 Estimation of Total soluble proteins, enzymatic antioxidants and MDA:

The bovine serum albumin was employed as a standard for the analysis of total soluble proteins (TSP) (Bradford, 1976). According to a known method, a protein-dye complex was prepared using 5 µL of supernatant, 95 µL of 150 mM NaCl and 1 mL of dye reagent, and then analyzed at 595 nm.

The 40 mM H₂O₂, 50 mM potassium phosphate buffer (pH 7.0) and 20 mM guaiacol were mixed with 0.1 mL enzyme extract. One unit of enzyme activity expressed on a fresh weight basis corresponds to a change in absorbance at 0.01 units/min. The POD activity corresponds to a rise in absorbance at 470 nm, while a decrease in absorbance at 240 nm provides CAT activity (Chance & Maehly, 1955).

The superoxide dismutase (SOD) activity was estimated from photochemical reduction of nitroblue tetrazolium (NBT) as a reference. The amount of enzyme that causes 50% inhibition of photochemical reduction of NBT corresponds to the one unit of SOD activity (Giannopolitis & Ries, 1977). The mixture of 50 µL enzyme extract, 13 mM methionine, 75 nM EDTA, 50 mM potassium phosphate buffer (pH 7.8), and 1.3 µM riboflavin was mixed with 50 µM NBT. Subsequently, the absorbance at 560 nm after irradiating the mixture under a fluorescent lamp.

The lipid peroxidation was estimated from the malondialdehyde (MDA) contents, measured using thiobarbituric acid (TBA). The 0.2 g seedling sample was mixed with 0.1% TCA and subjected

to centrifugation for 20 min. In the subsequent step, the supernatant was mixed with 0.05% TBA and 20% TCA and heated at 100 °C before recording the absorbance at 532 nm. The extinction coefficient ($155 \text{ mM}^{-1} \text{ cm}^{-1}$) was used to determine MDA (Heath & Packer, 1968).

2.3.2 Estimation of hydrolytic enzymes:

The TCA was used to measure protease activity following the casein digestion assay. The enzyme activity was defined as the amount of enzyme which released acid soluble fragments corresponding to 0.001 A₂₈₀ per minute at 37°C and pH 7.8. Enzymes activity was expressed on fresh weight basis (Drapeau, 1977). The mixture of 100 µL of enzyme extract, 2 mL of TCA and 2 mL of 1% casein was incubated, followed by absorbance measurement at 280 nm.

The α -amylase activity was measured according to the well-established methods using DNS reagent. The amount of amylase enzyme that will release 1 mg of maltose per minute under the established assay conditions was defined as 1 unit of enzyme activity. Enzymes activity was expressed on fresh weight basis (Varavinit et al., 2002). The 1 mL of enzyme extract, 1 mL of 1% starch solution and 1 mL DNS reagent were incubated for absorbance measurement at 540 nm.

2.3.3 Estimation of sugars:

The dinitrosalicylic acid was used for measuring of reducing sugars in seedling samples (Miller, 1959). Following the method, with 0.2 mL of extract in 1.8 mL dH₂O was mixed 1 mL DNS reagent and boiled for 10 min. Subsequently, the absorbance was recorded at 540 nm after adding 9 mL of distilled water. The phenol-sulfuric acid method was followed for the estimation of total sugars (Dubois et al., 1951). The 1 ml of extract was mixed with 0.5 ml of 5 % phenol and 2.5 ml of H₂SO₄ before recording the absorbance at 490 nm. Further, a standard curve of glucose was used for the determination of reducing and total sugars. Furthermore, the non-reducing sugars were calculated from the difference between reducing sugars and total sugars.

3. Results

The nanoprimed, hydroprimed and non-primed controls were germinated under normal and Cd stress conditions. The seedlings that originated from these seeds were used for biochemical analyses and compared with the control. The significance of the results at $p \leq 0.05$ and $p \leq 0.01$ was evaluated employing a two-way analysis of variance (ANOVA; factors: priming treatment [8 levels] and cadmium stress condition [2 levels: normal and stress] followed by Tukey's HSD post-hoc test using XLSTAT software. Values of three replicates with S.D and relative percentage have been presented in the Tables 1-3. The relative percentage difference (RPD) was calculated by the following formula:

$$\text{Relative Percentage Difference (RPD)}: \frac{(\text{Treatment} - \text{Control})}{\left(\frac{\text{Treatment} + \text{Control}}{2}\right)} \times 100$$

Baseline under normal condition was the corresponding non-stress control; baseline under Cd-stress condition was the corresponding Cd-stress control.

3.1 Effect of nanopriming on biochemical parameters in wheat seedlings:

The measured biochemical response was improved by the application of nanopriming under both non-stress and Cd-stress conditions, and the responses were found to be increasing with the increasing concentration of the nanoparticles. Overall, the responses were stronger in CS-SiO₂NP treatments (BT₁-BT₃) compared to the SiO₂NP treatments (AT₁-AT₃) in the presence of Cd stress.

In the presence of nanopriming, the total soluble protein (TSP) was significantly elevated in both conditions with generally a concentration-dependent response. Under control conditions, TSP was observed to be increased by 21-42% and by 23-37% with SiO₂NP (AT₁-AT₃) and CS-SiO₂NP (BT₁-BT₃), respectively. The second increase was 31-33% and 36-51% for the Cd stress. The higher response of the treatments coated with CS (especially BT₃) suggests higher effect of coating under Cd stress (**Table 1**).

Table 1. Effect of Chitosan-SiO₂NPs priming on enzymatic antioxidants and lipid peroxidation in wheat seedlings under normal and**Cadmium stress**

Treatments	Enzymatic Antioxidants									
	Normal					Under cadmium stress (5 μ M)				
	Total soluble proteins (TSP) (mg/g f. wt.)	Catalase (CAT) (units/g f.wt)	Peroxidase (POD) (units/g f.wt)	Superoxide dismutase (SOD) (units/g f.wt)	MDA (μ M/g f. wt)	Total soluble proteins (TSP) (mg/g f. wt.)	Catalase (CAT) (units/g f.wt)	Peroxidase (POD) (units/g f.wt)	Superoxide dismutase (SOD) (units/g f.wt)	MDA (μ M/g f. wt)
C₀ (Non-primed seeds)	202.167 $\pm$ 1.033 h	63 $\pm$ 1.082 b	17046.9 $\pm$ 1.9 c	136.097 $\pm$ 1.903 c	963.14 $\pm$ 3.855 ab	208 $\pm$ 1.022 h	38 $\pm$ 1.606 d	15098 $\pm$ 1 d	107.917 $\pm$ 1.083 d	997.790 $\pm$ 1.581 a
C_H	207.000 $\pm$ 1.026 h	64 $\pm$ 1.513 b	17087.3 $\pm$ 1.7 c (0.23%)	138.076 $\pm$ 1.924 c	903.21 $\pm$ 2.774 c	215.833 $\pm$ 1.167 h	39 $\pm$ 1.429 d	15435.8 $\pm$ 1.2 d	114.697 $\pm$ 1.303d	984.419 $\pm$ 1.742 a

(Hydro-primed)	(2.36 %)	(1.57%)		(1.5%)	(-6.43%)	(3.77%)	(2.60%)	(2.21%)	(5.40 %)	(-1.349%)
Sample AT₁	251.333±1.667	82±	19630.2±1.8	144.915±	895.33±	287± 1 e	58±	19102.5±	134.936±	941.145
(0.5ppm SiO₂NPs)	g (21.68%)	0.882 a (26.21%)	bc (14.08%)	0.8 b (6.40%)	1.661 c (-7.32%)	(31.92%)	1.333 c (41.67%)	1.5 bc (23.42%)	1.064 c (22.23%)	±1.855 b (-5.84%)
Sample AT₂	306.000±1.000	83±	24895±	150.364±	876.25±	293.167±	59±	21165.6±	134.426±	930.048
(1ppm SiO₂NPs)	c (40.86%)	0.208 a (27.40%)	1 a (37.42%)	1 a (9.79%)	1.742 d (-9.46%)	0.833 e (33.93%)	1.155 c (43.30%)	1.4 b (33.46%)	1.574 c (21.48%)	±1.952 b (-7.03%)
Sample AT₃	311.500±1.500	83±	25796.1±	152.525±	849.01±	291.833±	61±	25866.2±	138.864±	925.452
(1.5ppm SiO₂NPs)	c (42.57%)	1.028 a (27.40%)	1 .39 a (40.84%)	0.9 a (11.43%)	1.984 de (-12.58%)	1.167 e (33.6%)	1.764 c (46.47%)	1.2 a (52.57%)	1.136 c (25.10%)	±1.855 b (-7.52%)
Sample BT₁	254.833±0.767	82±	19744.7±	146.320±	889.98±	301± 1 d	54±	19094.5±1.5bc	142.740±	881.597
(0.5ppm Chitosan-SiO₂NPs)	g (23.05%)	1.528 a (26.21%)	1.7 bc (14.67%)	1.7 b (7.31%)	1.016 c (- 7.88%)	(36.54 %)	0.882 c (34.78%)	(23.37%)	1.26 b (28%)	±1.548 c (-12.36%)

Sample BT₂	277.333±0.667 f (31.35%)	81± 1.528 a (25%)	21784.1± 1.9 b (24.40%)	147.311± 1.4 b (7.98%)	868.17± 1.823 d (-10.38%)	326.5± 0.5 b (44.34%)	56± 0.882 c (38.30%)	21011.8± 1.2 b (32.75%)	144.926± 1.074 b (29.27%)	870.145 ±1.403 d (-13.67%)
Sample BT₃	296.500±1.5 d (37.83%)	84± 1.155 a (28.57%)	24830.4± 1.6 a (40.08%)	152.550± 1.6 a (11.42%)	815.09± 0.903 f (-16.65%)	353± 0.985 a (51.70 %)	63± 1.028 b (49.50%)	25120.6± 1.4 a (49.84%)	151.266± 1.734 a (33.45%)	815.532 ±1.468 f (-20.10%)
<i>Note: The percentage values were calculated based on the non-stress control and Cd-stress control, respectively.</i>										

Nanoprimering had a general positive effect on the expression of antioxidant enzymes. Under normal conditions, CAT activity was elevated by 25-28% in both SiO₂NPs and CS-SiO₂NPs while under Cd stress there was a higher elevation of CAT activity, 46% for AT₃ and 49% for BT₃. Activity of POD also increased with the concentration of the nanoparticles from 14-40% under normal conditions and 23-52% under Cd stress, with maximum activity at AT₃ and BT₃, respectively. The SOD activity had relatively small increases under the normal condition (6-11%) but the activity was greater under Cd stress (25% for AT₃ and 33% for BT₃). In general, the antioxidant response was stronger under Cd stress, especially at the highest concentration of CS-SiO₂NP (**Table 1**).

However, the MDA content decreased with increasing concentrations of nanoparticles, showing a progressive decrease in lipid peroxidation, whereas the antioxidant enzymes increased with nanoparticle concentration. When no stress was applied the change in MDA was 7-12% for SiO₂NPs and 7-16% for CS-SiO₂NPs. In Cd stress, the reductions were 5-7% and 12-20% respectively, the most considerable decrease being for BT₃. Therefore, CS coating improved (under Cd stress) the reduction of lipid peroxidation (**Table 1**).

Nanoprimering has also increased the activities of hydrolytic enzymes, but the sensitivity of the enzyme to nanoprimering varied among the treatments and stress conditions. In normal conditions, the activity of α -amylase increased gradually from 28% for AT₁ to 70% for AT₃, and from 33% for BT₁ to 93% for BT₃ in the CS-coated treatments. The activity of α -amylase under Cd stress was enhanced by 31-53% by SiO₂NPs and 24-51% by CS-SiO₂NPs. In control conditions, protease activity also increased in a concentration-dependent manner, with AT₁-AT₃ concentration, the activity increased by 5-25% and BT₁-BT₃ concentration, 6-34%. The increases were 33% for AT₃ and 26% for BT₃ under Cd stress. The results showed a significant rise in the activity of α -amylase and protease especially at the high concentration of nanoparticles after nanoprimering, suggesting an increase in hydrolytic activity after nanoprimering (**Table 2**).

Table 2. Effect of Chitosan-SiO₂NPs priming on hydrolytic enzymes in wheat seedlings under normal and Cadmium stress

Treatments	Hydrolytic Enzymes			
	Normal		Under cadmium stress (5 µM)	
	α- Amylase (units/g f. wt.)	Protease activity (Units/g f. wt.)	α- Amylase (units/g f. wt.)	Protease activity (Units/g f. wt.)
C₀ (Non-primed seeds)	8.075± 0.187 i	2678.5± 0.45 h	12.104± 0.032 g	2587± 0.671 h
C_H (Hydro-primed)	8.642± 0.106 i (6.78%)	2754.5± 0.5 g (2.79%)	14.179± 0.020 f (15.79%)	2710.5± 0.326 g (4.64%)
Sample AT₁ (0.5ppm SiO₂NPs)	10.708± 0.292 h (28.03%)	2822± 2.5 f (5.24%)	16.547 ± 0.045 d (31.01%)	2837.5± 1.5 f (9.22%)
Sample AT₂ (1ppm SiO₂NPs)	12.132± 0.187 g (40.15%)	2922.5± 1.45 e (8.71%)	19.057± 0.094 c (44.62%)	2995.5± 0.5 e (14.62%)
Sample AT₃ (1.5ppm SiO₂NPs)	16.830± 0.170 d (70.31%)	3467.5± 1.259 c (25.68%)	21± 0.059 b (53.74%)	3612.5± 2.5 b (33.07%)
Sample BT₁	11.321± 0.068 h	2854.5± 1.05 f	15.453± 0.055 d	2932.5± 2.5 e

(0.5ppm Chitosan-SiO₂NPs)	(33.47%)	(6.36%)	(24.30%)	(12.50%)
Sample BT₂	16.519± 0.481 d	2992± 1.002 e	16.236± 0.064 d	3212.5± 1.5 d
(1ppm Chitosan-SiO₂NPs)	(71.19%)	(11.07%)	(29.16%)	(21.55%)
Sample BT₃	22.377 ± 0.062 a	3790.5± 0.5 a	20.519± 0.481 b	3372.5± 0.986 c
(1.5ppm Chitosan-SiO₂NPs)	(93.93%)	(34.38%)	(51.59%)	(26.34%)
<i>Note: The percentage values were calculated based on the non-stress control and Cd-stress control, respectively.</i>				

Nanopriming also influenced the carbohydrate metabolism, with the concentration of the nanoparticles generally increasing the concentration of the non-reducing sugars and total sugars. In normal conditions, the level of non-reducing sugars increased by 11-16% and 18-32% in SiO₂NPs and CS-SiO₂NPs respectively, while in Cd stress it was increased by 10-24% and 5-21% respectively. Results showed that the total sugars concentration was elevated by 20-27% with SiO₂NPs and 43% with AT₃ under non-stress conditions while under Cd stress, the total sugars concentration increased by 39% with CS-SiO₂NPs. The total effect of the nanoparticles on sugar accumulation was greater with higher concentrations, although some parameters were affected more than others by the CS-coating effect (**Table 3**).

Table 3. Effect of Chitosan- SiO₂ NPs priming on sugar contents in wheat seedlings under normal and Cadmium stress

Treatments	Sugar contents					
	Normal			Under cadmium stress (5 µM)		
	Reducing Sugars (mg/g f. wt.)	Non-reducing Sugars (mg/g f. wt.)	Total Sugars (mg/g f. wt.)	Reducing Sugars (mg/g f. wt.)	Non-reducing Sugars (mg/g f. wt.)	Total Sugars (mg/g f. wt.)
C₀ (Non-primed seeds)	10.934± 0.054 f	13.227±0.079 h	24.160± 0.184 h	9± 0.246 g	14.422±0.058 f	23.422± 0.047 i
C_{II} (Hydro-primed)	11.976± 0.039 e (9.09%)	14.214±0.077 g (7.19%)	26.188± 0.225 g (8.05%)	10.853± 0.147 fg (18.67%)	14.650±0.085 f (1.57%)	25.504± 0.196 h (8.51%)
Sample AT₁ (0.5ppm SiO ₂ NPs)	14.795± 0.205 c (30.01%)	14.904±0.096 f (11.92%)	29.699± 0.3 ef (20.57%)	14.208± 0.092 c (44.88%)	15.973±0.093 d (10.20%)	30.181± 0.219 d (25.22%)
Sample AT₂ (1ppm SiO ₂ NPs)	15.286± 0.014 b (33.20%)	14.902±0.098 f (11.91%)	30.188± 0.312 e (22.18%)	14.491± 0.051 c (46.75%)	17.014±0.986 c (16.49%)	31.506± 0.494 d (29.43%)
Sample AT₃ (1.5ppm SiO ₂ NPs)	16.090± 0.010 ab (38.16%)	15.612±0.388 e (16.54%)	31.702± 0.298 d (27%)	18.024± 0.098 a (66.78%)	18.387±0.613 b (24.17%)	36.411± 0.589 b (43.42%)

Sample BT₁ (0.5ppm Chitosan-SiO₂NPs)	13.946± 0.054 cd (24.21%)	15.846±0.065 de (18.02%)	29.792± 0.106 f (20.88%)	13.090± 0.091 cd (37.03%)	15.216±0.284 e (5.36%)	28.306± 0.045 f (18.88%)
Sample BT₂ (1ppm Chitosan-SiO₂NPs)	15.110± 0.090 b (32.07%)	16.695±0.305 c (23.18%)	31.805± 0.03 d (27.32%)	15.141± 0.086 b (50.87%)	16.348±0.715 d (12.52%)	31.489± 0.076 d (29.38%)
Sample BT₃ (1.5ppm Chitosan-SiO₂NPs)	16.893± 0.107 ab (42.83%)	18.349±0.154 b (32.44%)	35.242± 0.261bc (37.31%)	17.039± 0.096 ab (61.74%)	17.952±0.548 c (21.81%)	34.991± 0.259 bc (39.61%)
<i>Note: The percentage values were calculated based on the non-stress control and Cd-stress control, respectively.</i>						

4. Discussion

Cd is a highly phytotoxic element that adversely affects physiological and biochemical parameters in germinating seeds, resulting in yield and quality losses. Therefore, seeds have developed various mechanisms to tolerate Cd and alleviate its toxicity.

Germinating seeds primarily use proteins as a source of C and N, vital for the synthesis of biomolecules during seedling establishment. An array of physiological and biochemical functions, including signal transduction, metabolic activities, and antioxidant defense, is controlled and coordinated by various enzymes, which are essentially proteins. Cd toxicity disrupts the TSP, generally disturbing vital processes, leading to abnormalities during germination and subsequent developmental stages (Bai et al., 2015). The Cd inactivates the enzymes either by replacing essential cofactors or by blocking thiol and carbonyl groups on their surface (Mostofa et al., 2015; Muszynska & Labudda, 2019; Rodriguez-Celma et al., 2010). Generally, seeds and developing plants increase their TSP content to counter the abiotic stresses, including HMs. The rice seedlings increased TSP and counterbalanced the toxic effects of Cd under SiO₂NPs treatments (Faisal et al., 2024). Spinach alleviated the Cd toxicity by increasing TSP following SiO₂NPs foliar application (Ullah et al., 2025). The exogenous application of CS remodeled biochemical parameters in wheat seedlings, including antioxidant enzyme activity, sugar levels, and increased TSP (17-36%) as a measure of Cd stress tolerance (Liu et al., 2021). The CS applications at lower doses upregulated the contents of antioxidants, chlorophyll, and TSP in pea (*Pisum sativum* L.) to alleviate Cd toxicity (Rasheed et al., 2020). The seed priming with CS activated the antioxidant machinery and increased TSP as a Cd acclamatory response in black seeds (*Nigella sativa* L.) (Arab et al., 2025). In a recent study, the foliar applications of CSNPs in *Solanum lycopersicum* protected against Cd toxicity by regulating enzymatic and non-enzymatic antioxidants, osmoprotectants and TSP contents (Faizan et al., 2021). The CS application controlled the glycolysis and protein expression as a stress-responsive measure in tomato seedlings (Attia et al., 2021). The CS increased the aspartate, glutamate and other amino acids, activated enzymes for N reduction, and modulated carbon and nitrogen assimilation in wheat

seedlings (Zhang et al., 2017). Accordingly, in our study, TSP content increased significantly following nanopriming with CS-coated SiO₂NPs, counteracting the negative impacts of Cd in wheat seedlings. The overexpression of TSP as key enzymes may have regulated antioxidant defense, metabolic activities, and cellular signaling during germination and seedling development.

Germinating seeds and growing plants activate antioxidant machinery as a main counter-step to resist HM's toxicity. They maintain controlled ROS production for the integrity of biomembranes by upregulating SOD, POD and CAT and neutralize HM toxicity (Moreira et al., 2020; Yang et al., 2012). Rice seedlings counteracted the Cd toxicity by upregulating its antioxidant potential due to SiO₂NPs treatments (Ashraf et al., 2024; Muhammad et al., 2025). The *Momordica charantia* L. seedlings activated the enzymatic antioxidants, showing significant resistance to Cd following nano-SiO₂ treatment (Sun et al., 2023). The *Satureja hortensis* L. also showed tolerance to toxic effects of Cd by upregulating CAT, SOD and POD activities following the foliar application of SiO₂NPs (Farajzadeh Memari-Tabrizi et al., 2021). The exogenous CS applications activated the enzymatic antioxidants and enhanced photosynthetic efficiency, enabling edible rape plants (*Brassica napus* L.) to withstand the detrimental effects of Cd stress (Zong et al., 2017). The CS treatments significantly improved antioxidant status and upregulated photosynthesis, protecting the maize (*Zea mays* L.) seedlings from Cd toxicity (Qu et al., 2019). In our case, the priming treatments activated the SOD, POD and CAT activities as a counterbalancing approach to resist Cd toxicity. The higher antioxidant activities were correlated with the lower level of lipid peroxidation, indicating that the antioxidants either prevent or slow the ROS mediated cellular damage. The significant decrease in MDA levels was correlated with nanopriming, indicating decrease in oxidative stress and Cd toxicity associated with increased antioxidant activity.

The hydrolytic enzymes, including α -amylase and protease, play a vital role in mobilizing stored reserves as a stress-acclamatory response during germination and plant growth. The hydrolytic enzymes exhibit reduced efficiency and adverse impacts due to Cd toxicity, leading to interrupted germination and seedling development (Anwar et al., 2021). The α -amylase loses its efficiency and

functioning after the replacement of Ca with Cd (Shiyu et al., 2020). The wheat seedlings exhibited resistance to Cd by modulating the biochemical attributes, including activation of hydrolytic enzymes, following SiO₂NPs treatments (Li et al., 2025). The maize seedlings also exhibited sustainable growth after SiO₂NPs treatments (Wang et al., 2024). The SiO₂NPs treatments of stevia positively influenced germination by activating the hydrolytic enzymes and other biochemical parameters (Hasanaklou et al., 2023). The upregulated hydrolytic enzymes induced tolerance against Cd toxicity following the foliar application of SiO₂NPs in wheat (Thind et al., 2021). Wheat seed priming with CS increased the activity of protease and α -amylase along with other biochemical features, resulting in improved germination and seedling development (Hameed et al., 2013). In this study, the protease and α -amylase exhibited increased activity, inhibiting Cd toxicity and ensuring sustainable seedling development.

Cd toxicity negatively influences and reduces sugar contents during seed germination and other developmental stages (Mishra et al., 2014). Seeds and growing plants use sugars as vital resources against the negative impacts of HMs. They provide insulation against HM and assist germination and seedling development by modulating biochemical attributes, respiration, and metabolic activities (Jeandet et al., 2022). They insulate the cellular structure by regulating ROS and acting as osmoprotectants (Farooq et al., 2022; O'Hara et al., 2013; Sami et al., 2016). Maize seedlings alleviated the toxic impacts of Cd by ROS scavenging role of sugars (Karalija & Selovic, 2018). Also, the rice seedlings mitigated Cd toxicity by regulating ROS with non-reducing sugars and regulation of metabolism (Wang et al., 2020). The wheat crop performed better with readjusted biochemistry and sugar contents following the silicon NP treatments (Li et al., 2023). In this study, priming treatments significantly increased the sugar contents as a countermeasure to resist Cd toxicity.

The significant improvements of the chitosan coated SiO₂NPs (BT₁, BT₂, BT₃) could be explained by the coating of chitosan, which provides both physicochemical and biological enhancement for the measured biochemical parameters that were consistently better than that of the

uncoated SiO₂NPs (AT₁, AT₂, AT₃). The positively charged seed surface carries a negative charge, and chitosan is a cationic biopolymer that contains free amino and hydroxyl groups, which can electrostatically interact with the seed surface, ensuring the retention of nanoparticles on the seed surface and improving the contact between the seed surface and the nanoparticles during imbibition at the initial stage (Hameed et al., 2013). The improved seed–particle interaction might help the Si to be absorbed into the plant and to alleviate Cd toxicity by improving the integrity of cell walls, antioxidant defense and reducing the bioavailability of Cd to the plant via co-precipitation and intracellular compartmentalization (Wang et al., 2022). Furthermore, chitosan has the ability to activate plant defense signaling pathways on its own, and influence expression of antioxidant-related genes, as well as biosynthesis of osmoprotectants (Mukarram et al., 2023). These synergistic effects would be stronger when coupled with the SiO₂NPs, offering twofold benefits: increased Si-mediated stress protection, decreased Cd bioavailability, in addition to chitosan-mediated defense activation and antioxidant effect. Such a synergistic effect could account for the improved performance of coated nanoparticles relative to the equivalent uncoated SiO₂NPs at the same concentration.

Principal component analysis (PCA) biplots were generated using OriginPro 2025b (OriginLab Corporation, Northampton, MA, USA) to evaluate the collective effects of seed priming treatments on biochemical parameters of wheat seedlings under normal and Cd-stressed conditions (Figure 3 a, b). The first two principal components explained 94.21% and 96.44% of the total variance under normal and Cd stress conditions, respectively, indicating strong inter-variable correlations and reliability of the PCA model.

Under normal conditions (Figure 3a), primed treatments, particularly CS-coated SiO₂NPs (BT₁-BT₃), were positively associated with total soluble proteins (TSP) and antioxidant enzymes (CAT, POD, SOD), demonstrating that nanoprimering induced a metabolically activated state even in the absence of stress. Hydro-primed and control seeds showed weaker associations with these biochemical traits, indicating limited physiological enhancement.




Figure 3. PCA analysis of CS-coated SiO₂NPs against Cd

Under cadmium stress (Figure 3b), a clear separation was observed between lipid peroxidation (MDA) and antioxidant/metabolic parameters. CS-coated SiO₂NPs-primed seedlings clustered closely with CAT, POD, SOD, and TSP, whereas MDA was oriented in the opposite direction, confirming that nanoprimering mitigated Cd-induced oxidative damage. The stronger grouping of antioxidant enzymes suggests a coordinated defense response triggered by CS-coated SiO₂NPs.

Overall, the PCA indicated correlative patterns, suggesting that CS-coated SiO₂NPs seed priming was associated with distinct biochemical profile that included increased antioxidant activities, higher protein and sugar levels and lower MDA concentrations under Cd stress. These patterns are correlations among the measured variables that do not necessarily imply a direct mechanism.

Wheat seed preconditioning with CS-coated SiO₂NPs triggered a biochemically activated seed state that is consistent with a priming response, contributing to Cd tolerance during seedling establishment. It regulated the enzymatic antioxidants, hydrolytic enzymes and other biomolecules to mitigate the Cd toxicity for sustainable growth.

5. Conclusions

Nanopriming has become a facile and innovative approach for managing stress and supporting climate-smart and sustainable plant growth. Seed preconditioning with CS-coated SiO₂NPs significantly increased total soluble proteins and sugars to counteract Cd toxicity. It avoided peroxidation and reduced MDA by controlling ROS production through activation of antioxidant enzymes. Further, it upregulated the protease and α -amylase enzymes which positively influenced the germination and seedling development. The findings showed that nanopriming with CS-coated SiO₂NPs successfully reduced the effects of Cd toxicity by modulating biochemical attributes for sustainable establishment of seedlings.

6. Future Recommendations:

This study has a number of limitations that should be considered. First, the direct quantification of Cd uptake in the seedling roots and shoots by ICP-MS or AAS was not conducted, which would give a definite proof of decreased Cd uptake and should be considered as a priority for future research. Second, the uptake of silicon into seedling tissues was not quantified; follow-up studies should confirm delivery of nanoparticles by ICP-OES or ICP-MS analysis. Third, no histochemical ROS localization procedures were performed by DAB or NBT staining nor were fluorescent ROS detection procedures performed in the present study, and the MDA levels and SOD activity data are functional measurements of ROS within the scope of the present study. Fourth, gene expression analysis of antioxidant and stress responsive genes (SOD, POD, CAT, APX) was not feasible within the scope of the present work and it is recommended as an important step to provide molecular level proof of the observed biochemical responses. Fifth, the study was done only in the seedling stage, hence further testing of nanopriming efficacy in pot or soil condition is important for future research. Sixth, there was a lack of a chitosan-only control in the experimental design, which would provide further information regarding the independent contribution of the SiO₂NP core and the CS coating. The study of stability of the nanoparticles under the priming conditions (in aqueous suspension, $23 \pm 2^\circ\text{C}$ for 8h) and the release kinetics of the CS-coated SiO₂NPs during seed imbibition were not determined

and should be investigated in future studies. Seventh, the tested concentration range (0.5, 1.0 and 1.5 ppm) is only a preliminary proof of concept and needs to be optimized further under greenhouse conditions. Lastly, CS and amorphous SiO₂ are both considered as low-toxicity and biodegradable, but long-term environmental effects after multiple applications of nanoparticles, especially on soil microbiota and aquatic ecosystems, need to be evaluated in the future studies.

Acknowledgements

The authors acknowledge the Hi-Tech lab of the Government College University Faisalabad for provision of centrifugation facilities.

References

- Anonymous. (1990). Association of Official Seed Analysts (AOSA). Rules for testing seeds. *Seed Science & Technology*, 12, 1-112.
- Anwar, S., Shafiq, F., Nisa, Z.-u.-., Usman, U., Ashraf, M. Y., & Ali, N. (2021). Effect of cadmium stress on seed germination, plant growth and hydrolyzing enzymes activities in mungbean seedlings. *Journal of Seed Science*, 43, e202143042. doi:10.1590/2317-1545v43256006
- Arab, E., Ahangar, L., & Biabani, A. (2025). The Effect of Chitosan Priming on Germination and Biochemical Characteristics of Black Seed (*Nigella sativa* L.) under Cadmium Stress. *Journal of Crop Breeding*, 17(1), 142-158. doi:10.61186/jcb.17.1.142
- Ashraf, H., Ghouri, F., Liang, J., Xia, W., Zheng, Z., Shahid, M. Q., & Fu, X. (2024). Silicon dioxide nanoparticles-based amelioration of cd toxicity by regulating antioxidant activity and photosynthetic parameters in a line developed from wild rice. *Plants*, 13(12), 1715. doi:10.3390/plants13121715
- Aslam, M. M., Okal, E. J., & Waseem, M. (2023). Cadmium toxicity impacts plant growth and plant remediation strategies. *Plant Growth Regulation*, 99, 397-412. doi:10.1007/s10725-022-00917-7
- Attia, M. S., Osman, M. S., Mohamed, A. S., Mahgoub, H. A., Garada, M. O., Abdelmouty, E. S., & Abdel Latef, A. A. H. (2021). Impact of foliar application of chitosan dissolved in different organic acids on isozymes, protein patterns and physio-biochemical characteristics of tomato grown under salinity stress. *Plants*, 10(2), 388. doi:10.3390/plants10020388
- Bai, X., Dong, Y., Kong, J., Xu, L., & Liu, S. (2015). Effects of application of salicylic acid alleviates cadmium toxicity in perennial ryegrass. *Plant Growth Regulation*, 75, 695-706. doi:10.1007/s10725-014-9971-3

- Bhatia, P., & Gupta, M. (2022). Micronutrient seed priming: new insights in ameliorating heavy metal stress. *Environmental Science and Pollution Research*, 29(39), 58590-58606. doi:10.1007/s11356-022-21795-6
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-254. doi:10.1016/0003-2697(76)90527-3
- Chance, B., & Maehly, A. C. (1955). Assay of catalases and peroxidases. In *Methods in Enzymology* (Vol. 2, pp. 764-775): Academic Press.
- Ci, D., Jiang, D., Wollenweber, B., Dai, T., Jing, Q., & Cao, W. (2010). Cadmium stress in wheat seedlings: growth, cadmium accumulation and photosynthesis. *Acta Physiologiae Plantarum*, 32, 365-373. doi:10.1007/s11738-009-0414-0
- Drapeau, G. R. (1977). Cleavage at glutamic acid with *Staphylococcal protease*. In *Methods in Enzymology* (Vol. 47, pp. 189-191): Academic Press.
- Dubois, M., Gilles, K., Hamilton, J. K., Rebers, P. A., & Smith, F. (1951). A colorimetric method for the determination of sugars. *Nature*, 168, 167. doi:10.1038/168167a0
- Eevera, T., Kumaran, S., Djanaguiraman, M., Thirumaran, T., Le, Q. H., & Pugazhendhi, A. (2023). Unleashing the potential of nanoparticles on seed treatment and enhancement for sustainable farming. *Environmental Research*, 236, 116849. doi:10.1016/j.envres.2023.116849
- El Rasafi, T., Oukarroum, A., Haddioui, A., Song, H., Kwon, E. E., Bolan, N., Tack, F. M. G., Sebastian, A., Prasad, M. N. V., & Rinklebe, J. (2022). Cadmium stress in plants: A critical review of the effects, mechanisms, and tolerance strategies. *Critical Reviews in Environmental Science and Technology*, 52(5), 675-726. doi:10.1080/10643389.2020.1835435
- Faisal, M., Faizan, M., Soysal, S., & Alatar, A. A. (2024). Synergistic application of melatonin and silicon oxide nanoparticles modulates reactive oxygen species generation and the antioxidant defense system: a strategy for cadmium tolerance in rice. *Frontiers in Plant Science*, 15, 1484600. doi:10.3389/fpls.2024.1484600
- Faizan, M., Rajput, V. D., Al-Khuraif, A. A., Arshad, M., Minkina, T., Sushkova, S., & Yu, F. (2021). Effect of foliar fertigation of chitosan nanoparticles on cadmium accumulation and toxicity in *Solanum lycopersicum*. *Biology*, 10(7), 666. doi:10.3390/biology10070666
- Farajzadeh Memari-Tabrizi, E., Yousefpour-Dokhanieh, A., & Babashpour-Asl, M. (2021). Foliar-applied silicon nanoparticles mitigate cadmium stress through physio-chemical changes to improve growth, antioxidant capacity, and essential oil profile of summer savory (*Satureja hortensis* L.). *Plant Physiology and Biochemistry*, 165, 71-79. doi:10.1016/j.plaphy.2021.04.040

- Farooq, T., Nisa, Z. U., Hameed, A., Ahmed, T., & Hameed, A. (2022). Priming with copper-chitosan nanoparticles elicit tolerance against PEG-induced hyperosmotic stress and salinity in wheat. *BMC Chemistry*, 16, 23. doi:10.1186/s13065-022-00813-1
- Giannopolitis, C. N., & Ries, S. K. (1977). Superoxide dismutases: I. Occurrence in higher plants. *Plant Physiology*, 59(2), 309-314. doi:10.1104/pp.59.2.309
- Gill, R. A., Kanwar, M. K., Rodrigues dos Reis, A., & Ali, B. (2021). Editorial: Heavy metal toxicity in plants: Recent insights on physiological and molecular aspects. *Frontiers in Plant Science*, 12, 830682. doi:10.3389/fpls.2021.830682
- Hameed, A., Sheikh, M. A., Hameed, A., Farooq, T., Basra, S. M. A., & Jamil, A. (2013). Chitosan priming enhances the seed germination, antioxidants, hydrolytic enzymes, soluble proteins and sugars in wheat seeds. *Agrochimica*, 57(2), 31-46.
- Hasanaklou, N. T., Mohagheghi, V., Hasanaklou, H. T., Ma'mani, L., Malekmohammadi, M., Moradi, F., & Dalvand, Y. (2023). Seed nano-priming using silica nanoparticles: effects in seed germination and physiological properties of *Stevia rebaudiana* Bertoni. *Chemical and Biological Technologies in Agriculture*, 10, 96. doi:10.1186/s40538-023-00445-0
- Hassan, M. J., Raza, M. A., Rehman, S. U., Ansar, M., Gitari, H., Khan, I., Wajid, M., Ahmed, M., Shah, G. A., Peng, Y., & Li, Z. (2020). Effect of cadmium toxicity on growth, oxidative damage, antioxidant defense system and cadmium accumulation in two sorghum cultivars. *Plants*, 9(11), 1575. doi:10.3390/plants9111575
- Heath, R. L., & Packer, L. (1968). Photoperoxidation in isolated chloroplasts: I. Kinetics and stoichiometry of fatty acid peroxidation. *Archives of Biochemistry and Biophysics*, 125(1), 189-198. doi:10.1016/0003-9861(68)90654-1
- Hou, D., O'Connor, D., Igalavithana, A. D., Alessi, D. S., Luo, J., Tsang, D. C., Sparks, D. L., Yamauchi, Y., Rinklebe, J., & Ok, Y. S. (2020). Metal contamination and bioremediation of agricultural soils for food safety and sustainability. *Nature Reviews Earth & Environment*, 1(7), 366-381.
- Huybrechts, M., Cuypers, A., Deckers, J., Iven, V., Vandionant, S., Jozefczak, M., & Hendrix, S. (2019). Cadmium and plant development: An agony from seed to seed. *International Journal of Molecular Sciences*, 20(16), 3971. doi:10.3390/ijms20163971
- Jeandet, P., Formela-Lubowska, M., Labudda, M., & Morkunas, I. (2022). The role of sugars in plant responses to stress and their regulatory function during development. *International Journal of Molecular Sciences*, 23(9), 5161. doi:10.3390/ijms23095161
- Karalija, E., & Selovic, A. (2018). The effect of hydro and proline seed priming on growth, proline and sugar content, and antioxidant activity of maize under cadmium stress. *Environmental Science and Pollution Research*, 25, 33370-33380. doi:10.1007/s11356-018-3220-7

- Khan, M. N., Fu, C., Li, J., Tao, Y., Li, Y., Hu, J., Chen, L., Khan, Z., Wu, H., & Li, Z. (2023). Seed nanoprimering: How do nanomaterials improve seed tolerance to salinity and drought? *Chemosphere*, 310, 136911. doi:10.1016/j.chemosphere.2022.136911
- Kumar, N., Kumar, V., Bose, B., & Singhal, R. K. (2021). Cadmium toxicity in plants and alleviation through seed priming approach. *Plant Physiology Reports*, 26, 647–660. doi:10.1007/s40502-021-00619-8
- Li, K., Li, Y., Yin, X., Yuan, Y., Yin, J., Chen, Y., & Zhu, Y. (2025). Silica nanoparticles enhanced seed germination and seedling growth of drought-stressed wheat by modulating antioxidant enzymes and mitigating lipid peroxidation. *Environmental Science: Nano*, 12(6), 3231-3246. doi:10.1039/d5en00214a
- Li, Y., Xi, K., Liu, X., Han, S., Han, X., Li, G., Yang, L., Ma, D., Fang, Z., Gong, S., Yin, J., & Zhu, Y. (2023). Silica nanoparticles promote wheat growth by mediating hormones and sugar metabolism. *Journal of Nanobiotechnology*, 21, 2. doi:10.1186/s12951-022-01753-7
- Liu, H., Yang, L., Li, N., Zhou, C., Feng, H., Yang, J., & Han, X. (2020). Cadmium toxicity reduction in rice (*Oryza sativa* L.) through iron addition during primary reaction of photosynthesis. *Ecotoxicology and Environmental Safety*, 200, 110746. doi:10.1016/j.ecoenv.2020.110746
- Liu, J., Gai, L., & Zong, H. (2021). Foliage application of chitosan alleviates the adverse effects of cadmium stress in wheat seedlings (*Triticum aestivum* L.). *Plant Physiology and Biochemistry*, 164, 115-121. doi:10.1016/j.plaphy.2021.04.038
- Ma, J., Faqir, Y., Chai, Y., Wu, S., Luo, T., Liao, S., Kaleri, A. R., Tan, C., Qing, Y., Kalhor, M. T., Umer, N., & Hadir, W. (2023). Chitosan microspheres-based controlled release nitrogen fertilizers enhance the growth, antioxidant, and metabolite contents of Chinese cabbage. *Scientia Horticulturae*, 308, 111542. doi:10.1016/j.scienta.2022.111542
- Miller, G. L. (1959). Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Analytical Chemistry*, 31(3), 426-428. doi:10.1021/ac60147a030
- Mishra, B., Sangwan, R. S., Mishra, S., Jadaun, J. S., Sabir, F., & Sangwan, N. S. (2014). Effect of cadmium stress on inductive enzymatic and nonenzymatic responses of ROS and sugar metabolism in multiple shoot cultures of Ashwagandha (*Withania somnifera* Dunal). *Protoplasma*, 251, 1031-1045. doi:10.1007/s00709-014-0613-4
- Moreira, I. N., Martins, L. L., & Mourato, M. P. (2020). Effect of Cd, Cr, Cu, Mn, Ni, Pb and Zn on seed germination and seedling growth of two lettuce cultivars (*Lactuca sativa* L.). *Plant Physiology Reports*, 25, 347-358. doi:10.1007/s40502-020-00509-5
- Mostofa, M. G., Rahman, A., Ansary, M. M. U., Watanabe, A., Fujita, M., & Tran, L.-S. P. (2015). Hydrogen sulfide modulates cadmium-induced physiological and biochemical responses to alleviate cadmium toxicity in rice. *Scientific Reports*, 5, 14078. doi:10.1038/srep14078

-
- Muhammad, S., Ulhassan, Z., Munir, R., Yasin, M. U., Islam, F., Zhang, K., Chen, W., Jan, M., Afzal, M., Muhammad, A., Hannan, F., & Zhou, W. (2025). Nanosilica and salicylic acid synergistically regulate cadmium toxicity in rice. *Environmental Pollution*, 364, 125331. doi:10.1016/j.envpol.2024.125331
- Mukarram, M., Ali, J., Dadkhah-Aghdash, H., Kurjak, D., Kacik, F., & Durkovic, J. (2023). Chitosan-induced biotic stress tolerance and crosstalk with phytohormones, antioxidants, and other signalling molecules. *Frontiers in Plant Science*, 14, 1217822. doi:10.3389/fpls.2023.1217822
- Muszynska, E., & Labudda, M. (2019). Dual role of metallic trace elements in stress biology—from negative to beneficial impact on plants. *International Journal of Molecular Sciences*, 20(13), 3117. doi:10.3390/ijms20133117
- Nandhini, R., Rajeswari, E., Harish, S., Sivakumar, V., Gangai Selvi, R., & Jaya sundrasharmila, D. (2025). Role of chitosan nanoparticles in sustainable plant disease management. *Journal of Nanoparticle Research*, 27, 13. doi:10.1007/s11051-024-06203-z
- Nile, S. H., Thiruvengadam, M., Wang, Y., Samynathan, R., Shariati, M. A., Rebezov, M., Nile, A., Sun, M., Venkidasamy, B., Xiao, J., & Kai, G. (2022). Nano-priming as emerging seed priming technology for sustainable agriculture—recent developments and future perspectives. *Journal of Nanobiotechnology*, 20, 254. doi:10.1186/s12951-022-01423-8
- O'Hara, L. E., Paul, M. J., & Wingler, A. (2013). How do sugars regulate plant growth and development? New insight into the role of trehalose-6-phosphate. *Molecular Plant*, 6(2), 261-274.
- Qu, D. Y., Gu, W. R., Zhang, L. G., Li, C. F., Chen, X. C., Li, J., Li, L. J., Xie, T. L., & Wei, S. (2019). Role of chitosan in the regulation of the growth, antioxidant system and photosynthetic characteristics of maize seedlings under cadmium stress. *Russian Journal of Plant Physiology*, 66, 140-151. doi:10.1134/S102144371901014X
- Rasheed, R., Ashraf, M. A., Arshad, A., Iqbal, M., & Hussain, I. (2020). Interactive effects of chitosan and cadmium on growth, secondary metabolism, oxidative defense, and element uptake in pea (*Pisum sativum* L.). *Arabian Journal of Geosciences*, 13, 847. doi:10.1007/s12517-020-05871-0
- Rashid, A., Schutte, B. J., Ulery, A., Deyholos, M. K., Sanogo, S., Lehnhoff, E. A., & Beck, L. (2023). Heavy metal contamination in agricultural soil: Environmental pollutants affecting crop health. *Agronomy*, 13(6), 1521. doi:10.3390/agronomy13061521
- Rodriguez-Celma, J., Rellan-Alvarez, R., Abadia, A., Abadia, J., & Lopez-Millan, A.-F. (2010). Changes induced by two levels of cadmium toxicity in the 2-DE protein profile of tomato roots. *Journal of Proteomics*, 73(9), 1694-1706. doi:10.1016/j.jprot.2010.05.001

-
- Saberi Riseh, R., Vatankhah, M., Hassanisaadi, M., & Varma, R. S. (2024). A review of chitosan nanoparticles: Nature's gift for transforming agriculture through smart and effective delivery mechanisms. *International Journal of Biological Macromolecules*, 260(2), 129522. doi:10.1016/j.ijbiomac.2024.129522
- Saleh, R. I., Kim, M., Baek, S. Y., & Cha, C. (2022). Chitosan-functionalized silica nanoparticles as a multifunctional coating material for improved water repellency, antimicrobial activity and mechanical strength of degradable bioplastics. *Cellulose*, 29, 7691-7701. doi:10.1007/s10570-022-04745-w
- Sami, F., Yusuf, M., Faizan, M., Faraz, A., & Hayat, S. (2016). Role of sugars under abiotic stress. *Plant Physiology and Biochemistry*, 109, 54-61. doi:10.1016/j.plaphy.2016.09.005
- Sarkar, M. R., Rashid, M. H.-o., Rahman, A., Kafi, M. A., Hosen, M. I., Rahman, M. S., & Khan, M. N. (2022). Recent advances in nanomaterials based sustainable agriculture: An overview. *Environmental Nanotechnology, Monitoring and Management*, 18, 100687. doi:10.1016/j.enmm.2022.100687
- Shiyu, Q., Hongen, L., Zhaojun, N., Rengel, Z., Wei, G., Chang, L., & Peng, Z. (2020). Toxicity of cadmium and its competition with mineral nutrients for uptake by plants: A review. *Pedosphere*, 30(2), 168-180.
- Singh, P. M., Tiwari, A., Maity, D., & Saha, S. (2022). Recent progress of nanomaterials in sustainable agricultural applications. *Journal of Materials Science*, 57, 10836-10862. doi:10.1007/s10853-022-07259-9
- Sun, H., He, S., Liu, T., Zhang, Q., Yu, J., Gao, Y., & Wang, X. (2023). Alleviation of cadmium toxicity by nano-silicon dioxide in *Momordica charantia* L. seedlings. *Journal of Soil Science and Plant Nutrition*, 23(1), 1060-1069. doi:10.1007/s42729-022-01103-4
- Suwanchaikasem, P., Idnurm, A., Selby-Pham, J., Walker, R., & Boughton, B. A. (2024). The impacts of chitosan on plant root systems and its potential to be used for controlling fungal diseases in agriculture. *Journal of Plant Growth Regulation*, 43, 3424–3445. doi:10.1007/s00344-024-11356-1
- Thind, S., Hussain, I., Rasheed, R., Ashraf, M. A., Perveen, A., Ditta, A., Hussain, S., Khalil, N., Ullah, Z., & Mahmood, Q. (2021). Alleviation of cadmium stress by silicon nanoparticles during different phenological stages of Ujala wheat variety. *Arabian Journal of Geosciences*, 14, 1028. doi:10.1007/s12517-021-07384-w
- Ullah, H., Sheng, Y., Wang, W., Wang, Z., Yang, H., & Dobbie, S. (2025). Application of SiO₂ nanoparticles to address CdS NPs contamination in spinach. *Plant Stress*, 16, 100811. doi:10.1016/j.stress.2025.100811

584 Varavinit, S., Chaokasem, N., & Shobsngob, S. (2002). Immobilization of a thermostable alpha-
585 amylase. *Science Asia*, 28(3), 247-251. doi:10.2306/scienceasia1513-1874.2002.28.247

586 Wang, K., Li, F., Gao, M., Huang, Y., & Song, Z. (2020). Mechanisms of trehalose-mediated
587 mitigation of Cd toxicity in rice seedlings. *Journal of Cleaner Production*, 267, 121982.
588 doi:10.1016/j.jclepro.2020.121982

589 Wang, L., Ning, C., Pan, T., & Cai, K. (2022). Role of silica nanoparticles in abiotic and biotic stress
590 tolerance in plants: A review. *International Journal of Molecular Sciences*, 23(4), 1947.
591 doi:10.3390/ijms23041947

592 Wang, M., Sun, G., Li, G., Hu, G., Fu, L., Hu, S., Yang, J., Wang, Z., & Gu, W. (2024). Effects of
593 multi walled carbon nanotubes and nano-SiO₂ on key enzymes for seed germination and
594 endogenous hormone level in maize seedling. *Agronomy*, 14(12), 2908.
595 doi:10.3390/agronomy14122908

596 Weisany, W., Razmi, J., Eshaghabadi, A. H., & Pashang, D. (2024). Silicon nanoparticles (SiNP): A
597 novel and sustainable strategy for mitigating environmental stresses in plants. *Journal of Soil
598 Science and Plant Nutrition*, 24, 2167-2191. doi:10.1007/s42729-024-01790-1

599 Wiszniewska, A. (2021). Priming strategies for benefiting plant performance under toxic trace metal
600 exposure. *Plants*, 10(4), 623. doi:10.3390/plants10040623

601 Wojtyła, L., Lechowska, K., Kubala, S., & Garnczarska, M. (2016). Molecular processes induced in
602 primed seeds—increasing the potential to stabilize crop yields under drought conditions.
603 *Journal of Plant Physiology*, 203, 116-126. doi:10.1016/j.jplph.2016.04.008

604 Yang, S., Xie, J., & Li, Q. (2012). Oxidative response and antioxidative mechanism in germinating
605 soybean seeds exposed to cadmium. *International Journal of Environmental Research and
606 Public Health*, 9(8), 2827-2838. doi:10.3390/ijerph9082827

607 Zhang, H., Xu, Z., Guo, K., Huo, Y., He, G., Sun, H., Guan, Y., Xu, N., Yang, W., & Sun, G. (2020).
608 Toxic effects of heavy metal Cd and Zn on chlorophyll, carotenoid metabolism and
609 photosynthetic function in tobacco leaves revealed by physiological and proteomics analysis.
610 *Ecotoxicology and Environmental Safety*, 202, 110856. doi:10.1016/j.ecoenv.2020.110856

611 Zhang, X., Li, K., Xing, R., Liu, S., & Li, P. (2017). Metabolite profiling of wheat seedlings induced
612 by chitosan: Revelation of the enhanced carbon and nitrogen metabolism. *Frontiers in Plant
613 Science*, 8, 2017. doi:10.3389/fpls.2017.02017

614 Zhu, T., Li, L., Duan, Q., Liu, X., & Chen, M. (2021). Progress in our understanding of plant
615 responses to the stress of heavy metal cadmium. *Plant Signaling and Behavior*, 16(1),
616 1836884. doi:10.1080/15592324.2020.1836884

617 Zong, H., Liu, S., Xing, R., Chen, X., & Li, P. (2017). Protective effect of chitosan on photosynthesis
618 and antioxidative defense system in edible rape (*Brassica rapa* L.) in the presence of

List of Figures

Figure 1. Preparation of CS-coated SiO₂NPs as nanoprimer agent

Figure 2. Characterization of CS-coated SiO₂NPs

Figure 3. PCA analysis of CS-coated SiO₂NPs against Cd

List of Tables

Table 1: Effect of Chitosan-SiO₂NPs priming on enzymatic antioxidants and lipid peroxidation in wheat seedlings under normal and Cadmium stress

Table 2: Effect of Chitosan-SiO₂NPs priming on hydrolytic enzymes in wheat seedlings under normal and Cadmium stress

Table 3: Effect of Chitosan-SiO₂NPs priming on sugar contents in wheat seedlings under normal and Cadmium stress

Declarations

Funding

There was no funding for this study.

Conflicts of Interest

“The authors declare no conflict of interest”.

Credit Author Statement

Arruje Hameed: Conceptualization, Writing, Supervision. **Ayesha Saeed:** Experimentation, data curation and analyses. **Razia Noreen:** Sources, Member of the supervisory committee. **Nosheen Aslam:** Sources, Member of the supervisory committee.